

REGULAR AND BIREGULAR MODULE ALGEBRAS

CHRISTIAN LOMP

ABSTRACT. Motivated by the study of von Neumann regular skew groups as carried out by Alfaro, Ara and del Rio in [1] we investigate regular and biregular Hopf module algebras. If A is an algebra with an action by an affine Hopf algebra H , then any H -stable left ideal of A is a direct summand if and only if A^H is regular and the invariance functor $(-)^H$ induces an equivalence of A^H -Mod to the Wisbauer category of A as $A\#H$ -module. Analogously we show a similar statement for the biregularity of A relative to H where A^H is replaced by $R = Z(A) \cap A^H$ using the module theory of A as a module over $A^e \bowtie H$ the envelopping Hopf algebroid of A and H . We show that every two-sided H -stable ideal of A is generated by a central H -invariant idempotent if and only if R is regular and A_m is H -simple for all maximal ideals m of R . Further sufficient conditions are given for $A\#H$ and A^H to be regular.

1. INTRODUCTION

Motivated by the study of von Neumann regular of skew group rings by Alfaro et al. in [1] and by the studies of the regularity of fix rings by Goursad et all in [9] we look at the regularity of Hopf module algebras, their smash products and their subrings of invariants. To achieve our goal we will work in the following more general setting:

Let k be a commutative ring. An extension $A \subseteq B$ of k -algebras is said to have an additional module structure if there exists a ring homomorphism $\Psi : B \rightarrow \text{End}_k(A)$ such that $\Psi(a) = L_a$ for all $a \in A$, where L_a denotes the left multiplication of a on A . Then A is a cyclic left B -module with B -action $b \cdot a := \Psi(b)(a)$ for all $b \in B, a \in A$. Moreover $\alpha : B \rightarrow A$ with $(b)\alpha = b \cdot 1$ is an epimorphism of left B -modules. Note that we will write homomorphisms oposite of scalars. Furthermore $\phi : \text{End}_B(A) \rightarrow A$ with $\phi(f) = (1)f$ defines a ring homomorphism whose image is denote by A^B . In particular

$$A^B = \{a \in A \mid \forall b \in B : b \cdot a = (b)\alpha\} = \{a \in A \mid \forall b \in B \ \forall a' \in A : b \cdot (a'a) = (b \cdot a')a\}.$$

Defining for any B -module M :

$$M^B = \{m \in M \mid \forall b \in B \ \forall a \in A : b \cdot (am) = (b \cdot a)m\}$$

one also has functorial isomorphisms

$$\text{Hom}_B(A, M) \longrightarrow M^B \quad f \mapsto (1)f$$

such that $\text{Hom}_B(A, -)$ and $(-)^B$ are isomorphic functors (see [14] for details). In the terminology of [3], B is an A -ring with a right grouplike character.

Examples of the described situation are abundant in the theory of Hopf algebra actions where a Hopf algebra H (or more general a weak Hopf algebra) acts on an algebra A and $A \subseteq B =$

Key words and phrases. Regular rings, biregular rings, Hopf actions, smash products, envelopping Hopf algebroid.

This work was partially supported by Centro de Matemática da Universidade do Porto (CMUP), financed by FCT (Portugal) through the programs POCTI (Programa Operacional Ciência, Tecnologia, Inovação) and POSI (Programa Operacional Sociedade da Informação), with national and European community structural funds.

$A \# H$ is an extension with additional module structure. This also includes group action and Lie actions. Further examples are given by the enveloping algebra $A \subseteq A^e$ or more generally by the enveloping Hopf algebroid $A^e \bowtie H$ as defined in [14] (see also [13] or [7]), k -algebras A with an involution $*$ with $B = A^e * G$ where $G = \langle \sigma \rangle$ is group generated by the automorphism σ of A^e defined by $\sigma(a \otimes b) = b^* \otimes a^*$ or certain extensions $A \subseteq B$ arising in the study of Banach algebras (see Cabrera et al. [2]).

In this paper we will characterize regular and biregular H -module algebras, generalising some known results on the regularity skew group rings.

All rings will be associative and unital. Ring homomorphisms are supposed to respect the unit. Throughout the text k will denote a commutative ring and A a k -algebra. We denote by $A^e := A \otimes A^{op}$ the enveloping algebra of A whose multiplication is defined as $(a \otimes b)(a' \otimes b') = aa' \otimes b'b$.

2. REGULAR MODULES

John von Neumann defined a ring R to be regular if for any element $a \in R$ there exists an element $b \in R$ such that $a = aba$. He showed in [17] that R is regular if and only if every cyclic (finitely generated) left (right) ideal of R is a direct summand. Later Auslander proved that the regularity of a ring can also be characterised by the property that any module is flat or equivalently that any submodule of a module is pure. Several authors have transferred the regularity condition to modules. A. Tuganbaev in [20] calls a left R -module M regular if any cyclic (finitely generated) submodule is a direct summand using the lattice theoretical approach, while J. Zelmanowitz in [24] followed the original elementwise definition of von Neumann and called a left R -module M regular if for any $m \in M$ there exists $f \in \text{Hom}_R(M, R)$ such that $(m)fm = m$.

The module theoretic version of Auslander's characterisation had been carried out by Fieldhouse [8] where he called a left R -module regular if any of its submodule is pure in the sense of P.M. Cohen. R. Wisbauer [21] used his ideas to define regularity for nonassociative rings (see also [22, Chapter 34]): Let R be an arbitrary ring and M a left R -module. The Wisbauer category $\sigma[M]$ is the subcategory of $R\text{-Mod}$ whose objects are the submodules of M -generated modules, i.e. submodules of factor modules of direct sums of copies of M . A module $P \in \sigma[M]$ is called finitely presented in $\sigma[M]$ if P is finitely generated and every exact sequence in $\sigma[M]$:

$$0 \longrightarrow K \longrightarrow L \longrightarrow P \longrightarrow 0$$

with L finitely generated implies K to be finitely generated. Note that P might be finitely presented in $\sigma[M]$ but not in $R\text{-Mod}$, for example take any simple module $P = M$. A short exact sequence in $\sigma[M]$ is called *pure* if any finitely presented module in $\sigma[M]$ is projective with respect to this sequence and a module $N \in \sigma[M]$ is called *flat* in $\sigma[M]$ if any short exact sequence

$$0 \longrightarrow K \longrightarrow L \longrightarrow N \longrightarrow 0$$

in $\sigma[M]$ is pure. Finally M is called *regular* if any module in $\sigma[M]$ is flat or equivalently if any short exact sequence in $\sigma[M]$ is pure.

2.1. Relative regularity. Let $A \subseteq B$ be an extension with additional module structure. Our first aim will be to characterise A as a regular B -module.

Proposition. *Let $A \subseteq B$ be an extension with additional module structure. The following statements are equivalent:*

- (a) *A is regular and finitely presented in $\sigma[B A]$;*

- (b) Every left B -stable ideal that is finitely generated as left B -module is a direct summand and A is finitely presented in $\sigma[B A]$.
- (c) A^B is von Neumann regular and A is a generator in $\sigma[B A]$.
- (d) A^B is von Neumann regular and $()^B = \text{Hom}_B(A, -) : \sigma[B A] \rightarrow A^B\text{-Mod}$ is an equivalence of categories.

In this case A is a projective generator in $\sigma[B A]$.

Proof. (a) $\Leftrightarrow$ (b) follows from [22, 37.4]

(a) $\Rightarrow$ (c) Since A is finitely presented and regular in $\sigma[B A]$, it is projective in $\sigma[B A]$. By As A is a cyclic B -module, by [22, 37.8], A is a (finitely generated) projective generator in $\sigma[B A]$ and A^B is regular.

(c) $\Leftrightarrow$ (d) is clear.

(c) $\Rightarrow$ (b) Since $B A$ is cyclic and a generator in $\sigma[B A]$ and since $A^B \simeq \text{End}_B(A)$ is regular and thus A is a faithfully flat A^B -module, we have by [22, 18.5(2)] that A is self-projective (and hence projective in $\sigma[B A]$). This implies that A is also finitely presented in $\sigma[B A]$. If U is a finitely generated B -stable left ideal of A , then $U = AI$ for $I = U^B$. U being finitely generated as B -module, implies I being finitely generated as right ideal of A^B . Thus $I = A^B e$ for some idempotent e and $U = AI = Ae$ is a direct summand of A , i.e. A is regular by [22, 37.4]. $\square$

2.2. Relative biregularity. If $A \subseteq B$ is an extension with additional modules structure $\Psi : B \rightarrow \text{End}_k(A)$, we might identify B with its image in $\text{End}_k(A)$ seeing it as an extension of the subalgebra generated by left multiplications of A . In order to study the two-sided B -stable ideals we might enlarge B by considering $B' = \langle B \cup M(A) \rangle \subseteq \text{End}_k(A)$. Note that all B -submodules of A are two-sided and $A^B \subseteq Z(A)$ if $M(A) \subseteq B \subseteq \text{End}_k(A)$. A B -stable ideal I is called prime if $JK \subseteq I$ implies $J \subseteq I$ or $K \subseteq I$ for any B -stable ideals J and K . I is semiprime if it is the intersection of prime B -stable ideals. A is B -semiprime if 0 is a prime as B -stable ideal or equivalently A does not contain any non-zero nilpotent B -stable ideal (see [13, 2.3]). If a cyclic B -stable ideal $B \cdot a$ is a direct summand of A , then there exists an idempotent $e \in A^B$ with $B \cdot a = B \cdot e = Ae$ and $A = Ae \oplus A(1 - e)$. A k -algebra A is called B -biregular if every cyclic B -stable ideal is a direct summand of A . In particular Proposition 2.1 applies to get a characterisation of B -biregular algebras A in case A is finitely presented in $\sigma[B A]$, namely that A is B -biregular if and only if A^B is a von Neumann regular ring with $(-)^B : \sigma[B A] \rightarrow A^B\text{-Mod}$ being an equivalence.

2.3. Properties of relative biregular algebras. In the next two subsections, we intend to characterise B -biregular algebras A without assuming that A is finitely presented in $\sigma[B A]$.

Proposition. Let $M(A) \subseteq B \subseteq \text{End}_k(A)$. Suppose that A is B -biregular. Then

- (1) A^B is von Neumann regular and A is B -semiprime.
- (2) A is a A^B -Ideal Algebra, i.e. the map $I \mapsto IA$ is a bijection between the ideals I of A^B and the B -stable ideals of A , whose inverse is given by $N \mapsto \text{Ann}_{A^B}(A/N) \simeq \text{Hom}_B(A/N, A)$.
- (3) Every finitely generated B -stable ideal of A is cyclic and is generated by some central idempotent in A^B .
- (4) For any B -stable ideal I of A , also A/I is B/I -biregular.
- (5) Every B -stable ideal of A is idempotent and equals the intersection of maximal B -stable ideals.

(6) *Every prime B -stable ideal is maximal.*

Proof. (1) Let $f \in \text{End}_B(A)$, then $(A)f = B(1)f$ is a direct summand in A by hypothesis, i.e. $(A)f = Ae$ with $e^2 = e \in A^B \subseteq Z(A)$. Since $A(1-e) \subseteq \text{Ker}(f) \subseteq l.\text{ann}((A)f) = A(1-e)$ also the kernel of f is a direct summand,. Hence by [23, 7.6], $\text{End}_B(A)$ and thus A^B is regular. Since no cyclic B -stable ideal is nilpotent, A is B -semiprime.

(2) A generates all cyclic B -stable ideals, i.e. ${}_B A$ is a self-generator and since A^B is regular by (1), ${}_B A$ is intern-projective by [23, 5.6]. Since A is a cyclic B -module, the claim then follows by [23, 5.9].

(3) Let Ae and Af be cyclic B -stable ideals with idempotents $e, f \in A^B$. Then $Ae + Af = A(e + f - ef) = A(e \uplus f)$, where $\uplus$ is the addition in the boolean ring of idempotents $B(A^B)$.

(4) By (2), every B -stable ideal I can be written as $I = JA$ with J ideal in $Z := A^B$. Hence the canonical projection $A = A \otimes_Z Z \rightarrow A/I \simeq A \otimes Z/J$ can be understood as the tensoring of the canonical projection of $Z \rightarrow Z/J$ by $A \otimes_Z -$, which respects direct sums.

(5) For every cyclic B -stable ideal $B \cdot x = Ae$ we have $(Ae)^2 = A^2 e^2 = Ae$. Hence $B \cdot x$ and thus any B -stable ideal is idempotent. Since there are no small B -submodules in A , we have $\text{Rad}({}_B A) = 0$ and 0 is the intersection of maximal B -stable ideals. By (4) we can use this argument to each A/I .

(6) Suppose A is B -prime and B -biregular. Let $0 \neq I = Ae$ be a cyclic B -stable ideal with idempotent e . As $A(1-e)$ is a B -stable ideal with $A(1-e)I = 0$, we have $A(1-e) = 0$, i.e. $I = A$ and A is B -simple. $\square$

2.4. Characterisation of relative biregularity. The next Proposition characterises biregular extensions $A \subseteq M(A) \subseteq B \subseteq \text{End}_k(A)$. Denote by $\text{Max}(A^B)$ the spectrum of maximal ideals of A^B and by A_m the localisation of A by a maximal ideal m of A^B . Note that if A^B is regular, then $A_m = A/mA$ by [23, 17.7] and in particular since mA is B -stable, we might consider $B \subseteq \text{End}_k(A/mA) = \text{End}_k(A_m)$. We say that A is B -simple if 0 and A are the only B -stable ideals of A .

Theorem. *The following statements are equivalent for an extension $M(A) \subseteq B \subseteq \text{End}_k(A)$.*

- (a) A is B -biregular;
- (b) A^B is regular and every maximal B -stable ideal M of A is of the form $M = AM^B$.
- (c) A^B is regular and A_m is B -simple for all $m \in \text{Max}(A^B)$.

Proof. (a) $\Rightarrow$ (b) the properties (i – iii) follow from Proposition 2.3 and (iv) follows from the fact if A is B -biregular then for any $x \in A : l.\text{ann}_A(Bx) = A(1-e)$ with $e^2 = e \in A^B$ is already a B -ideal.

(b) $\Rightarrow$ (c): Let m be a maximal ideal of A^B and let M be a maximal B -stable ideal containing $mA \subseteq M$. Since $M = M^B A$ we have

$$m \subseteq (mA)^B \subseteq M^B$$

which implies $M^B = m$ since $M \neq A$. Thus $mA = M$ and $A/M = A/mA = A_m$ is B -simple.

(c) $\Rightarrow$ (a) Let I be any B -stable ideal I of A . Then $I^B A \subseteq I$ and

$$(I^B A)_m = (I \cap A^B)_m A_m = (I_m \cap A_m^B) A_m.$$

If $I_m = A_m$ then $I_m^B = I_m \cap A_m^B = A_m^B$ and hence $(I^B A)_m = I_m$. If $I_m \neq A_m$, then $I_m = 0_m$ and therefore $I_m^B = 0_m$, i.e. $(I^B A)_m = I_m$. Since this holds for any maximal ideal m of A^B , we get $I = I^B A$ which shows that A is a self-generator as B -module.

□

2.5. Regular subring of invariants. Assume again that $A \subseteq B$ is any extension with additional module structure. In order to determine when the subring of invariants A^B is regular, we need first to borrow another notion from module theory.

Definition. A left R -module M is called *semi-projective*, if every diagram:

$$\begin{array}{ccc} & M & \\ & \downarrow g & \\ M & \xrightarrow{f} N & \longrightarrow 0 \end{array}$$

with $N \subseteq M$ can be completed by an endomorphism $h \in S := \text{End}_R(M)$ such that $hf = g$. As it is easily seen: M is semi-projective if and only if $\text{Hom}_R(M, Mf) = Sf$ for all $f \in S$.

Hence A is semi-projective as left B -module if $\forall x \in A^B : (Ax)^B = A^B x$.

Proposition. Let $A \subseteq B$ be an extension with additional module structure. Then A^B is von Neumann regular if and only if

- (1) A is semi-projective as left B -module and
- (2) every cyclic left ideal generated by an B -invariant element $x \in A^B$ is a direct summand of A as B -module.

Proof. If $A^B \simeq \text{End}_B(A)$ is regular, then ${}_B A$ is semi-projective by [23, 5.9]. Furthermore since the images of B -linear maps are direct summands and are precisely the cyclic B -stable left ideals generated by a B -invariant element we are done.

On the other hand assume that ${}_B A$ is semi-projective. Let $0 \neq x \in A^B$ then $B \cdot x = Ax$ is a direct summand of A as left B -module by hypothesis. Thus $A = Ax \oplus I$ as left B -modules. But then

$$A^B = (1)(\text{Hom}_B(A, Ax) \oplus \text{Hom}_B(A, I)) = (Ax)^B \oplus I^B = A^B x \oplus I^B.$$

Hence every cyclic left ideal of A^B is a direct summand, i.e. A^B is von Neumann regular. □

2.6. Large subring of invariants. If A is finitely presented and regular in $\sigma[{}_B A]$, then by Proposition 2.1 it is a projective generator in $\sigma[{}_B A]$. Weakening the generator condition J.Zelmanowitz called a left R -module M *retractable* if $\text{Hom}_R(M, N) \neq 0$ for all non-zero submodules $N \subseteq M$. For a module algebra extension $A \subseteq B$ we say that A^B is *large* in A if $I \cap A^B \neq 0$ for all B -stable left ideals of A or equivalently if A is a retractable B -module. A classical theorem of Bergmann and Isaacs says that if finite group G acts on an algebra A such that A is G -semiprime and has no $|G|$ -torsion, then R^G is large in R .

A purely module theoretical result by J.Zelmanowitz from [25] says now in our language:

Lemma. Let A be projective in $\sigma[{}_B A]$ and A^B large in A , then

- (1) If A^B is left self-injective, then A is a self-injective left B -module.
- (2) If A^B is von Neumann regular, then A is a non-singular in $\sigma[{}_B A]$, i.e. if $K \subseteq L$ is an essential extension in $\sigma[{}_B A]$, then $\text{Hom}_B(L/K, A) = 0$.

Proof. Zelmanowitz calls a left R -module M *fully retractable* if $\text{Hom}_R(M, N)g \neq 0$ for any $0 \neq g \in \text{Hom}_R(N, M)$ and submodule $N \subseteq M$. It is easy to see that self-projective retractable modules are fully retractable. Zelmanowitz proves in [25, Proposition on page 567] that M is self-injective

if M is fully retractable and left $\text{End}_R(M)$ self-injective. Property (2) follows from [25, Corollary on page 568]. $\square$

Note that a module M is non-singular in $\sigma[M]$ if and only if it is “polyform” in the sense of J. Zelmanowitz (see [23]).

2.7. As a consequence we have that if A is projective in $\sigma[B A]$ and A^B large in A , then A^B is regular and left self-injective if and only if A is injective and non-singular in $\sigma[B A]$, because the endomorphism ring of any self-injective polyform module is self-injective and regular by [23, 11.1].

3. RELATIVE SEMISIMPLE EXTENSIONS

Let $A \subseteq B$ be an extension of k -algebras. An element $c = \sum_i c_i \otimes c^i \in B \otimes_A B$ which is B -centralising, i.e. $bc = cb$ for all $b \in B$ is called a *Casimir element* for B over A (see [19] for the terminology). We say that a Casimir element acts unitarily on an element m of a left B -module M if $(\sum_i c_i c^i) \cdot m = m$.

Proposition. *Let $A \subseteq B$ be an extension with additional module structure and suppose that B has a Casimir element over A that acts unitarily on A , then the following hold:*

- (1) *c acts unitarily on any module in $\sigma[B A]$.*
- (2) *The k -linear map $\text{Hom}_A(M, N) \rightarrow \text{Hom}_B(M, N)$ with $f \mapsto \tilde{f} : [m \mapsto \sum c_i \cdot f(c^i \cdot m)]$ splits the embedding $\text{Hom}_B(M, N) \subseteq \text{Hom}_A(M, N)$ for any $N, M \in \sigma[B A]$.*

Proof. Let $\gamma := \sum c_i c^i$ and $\alpha : B \rightarrow A$ with $(b)\alpha = b \cdot 1$. Then α is left B -linear and $(a)\alpha = a$ for any $a \in A$. For all $a \in A$ we have $ac = \sum ac_i \otimes c^i = \sum c_i \otimes c^i a = ca$. Then also $(\sum ac_i c^i) \alpha = (\sum c_i c^i a) \alpha$ holds. Thus

$$(*) \quad a = a(\gamma)\alpha = \left(\sum ac_i c^i\right)\alpha = \left(\sum c_i c^i a\right)\alpha = \left(\sum c_i c^i\right) \cdot (a)\alpha = \gamma \cdot a.$$

(1) Let $M \in \sigma[B A]$. Then there exists a set Λ and a B -submodule $I \subseteq A^{(\Lambda)}$, such that M is isomorphic to a B -submodule of $A^{(\Lambda)}/I$. We identify M with a submodule of $A^{(\Lambda)}/I$. Let $m \in M$, then there are elements $a_\lambda \in A$ for $\lambda \in \Lambda$ such that $m = (a_\lambda)_\Lambda + I$. Now it follows with (*):

$$\gamma \cdot m = \gamma \cdot [(a_\lambda)_\Lambda + I] = (\gamma \cdot a_\lambda)_\Lambda + I = (a_\lambda)_\Lambda + I = m.$$

(2) Obviously $\tilde{f}$ is B -linear for all $f : M \rightarrow N$ since c is a Casimir element. If f was already B -linear, then using (1) we get for all $m \in M$:

$$\tilde{f}(m) = \sum c_i \cdot f(c^i \cdot m) = \left(\sum c_i c^i\right) \cdot f(m) = f(m),$$

i.e. $\tilde{f} = f$ showing that the embedding splits. $\square$

3.1. M is a (B, A) -semisimple B -module if any short exact sequence in $\sigma[B M]$ that splits as left A -module, also splits as left B -module (see [23, page 170]). Recall that Hirata and Sugano called a ring extension $A \subseteq B$ a semisimple extension if B is (B, A) -semisimple (see [10]).

Corollary. *If B has a Casimir element c which acts unitarily on A , then A is a (B, A) -semisimple B -module and for any $M \in \sigma[B A]$*

- *If M is N -projective as A -module for $N \in \sigma[B A]$, then M is also N -projective as B -module.*
- *If M is N -injective as A -module for $N \in B\text{-Mod}$, then M is also N -injective as B -module.*

In particular A is projective in $\sigma[B A]$.

Proof. Let $\pi : M \longrightarrow N$ be a projection in $\sigma[_B A]$ with $\pi(n) = n$. Then for any $n \in N$:

$$\tilde{\pi}(n) = \sum c_i \cdot \pi(c^i \cdot n) = \left(\sum c_i c^i \right) \cdot \pi(n) = \pi(n) = n.$$

Thus $\tilde{\pi}$ splits the embedding of N into M as B -module. In the same way one proves the statements (1). For (2) note that if $f : U \rightarrow M$ is B -linear, where U is a B -submodule of N , then there exists an A -linear map $g : N \rightarrow M$ such that $g|_U = f$. Set as before $\tilde{g} : N \rightarrow M$ which is B -linear. Then $\tilde{g}(u) = (\sum c_i c^i) \cdot f(u) = f(u)$. $\square$

3.2. In [10] Hirata and Sugano called a ring extension $A \subseteq B$ *separable* if there exists a Casimir element $c = \sum_i c_i \otimes c^i$ such that $\sum_i c_i c^i = 1$.

Corollary. *Let $A \subseteq B$ be an extension with additional module structure such that there exists a Casimir element in B which acts unitarily on A , then*

- (1) *If A is a semisimple artinian ring, then A is semisimple B -module.*
- (2) *If A is von Neumann regular and ${}_A B$ is finitely generated, then*
 - *A is a regular module in $\sigma[_B A]$;*
 - *A^B is a regular ring and*
 - *$(-)^B$ defines a Morita equivalence between $A^B\text{-Mod}$ and $\sigma[_B A]$.*
- (3) *If $\sigma[_B A] = B\text{-Mod}$, then $A \subseteq B$ is a semisimple extension.*

Proof. (1) Is clear since A is (B, A) -semisimple.

(2) Since ${}_A B$ is finitely generated, A is finitely presented in $\sigma[_B A]$. If $B \cdot a$ is a cyclic B -submodule of A , then by hypothesis $B \cdot a$ is also finitely generated as left A -module and hence a direct summand of A as left A -module. Thus $B \cdot a$ is also a direct summand of A as left B -module since A is (B, A) -semisimple. By 2.1 A is a regular module in $\sigma[_B A]$. Also by 2.1 we have that $\text{End}_B(A) \simeq A^B$ is regular and A is a progenerator in $\sigma[_B A]$ with equivalence $\text{Hom}_B(A, -) \simeq (-)^B : \sigma[_B A] \longrightarrow \text{End}_B(A) \simeq A^B$.

(3) If A is a subgenerator in $B\text{-Mod}$, then $B \in \sigma[_B A]$ is itself (B, A) -semisimple. $\square$

4. APPLICATIONS TO HOPF ALGEBRA ACTIONS

Let H be a Hopf algebra over k acting on an algebra A , i.e. A is a left H -module algebra. The smash product of A and H is denoted by $A \# H$ whose underlying k -module is $A \otimes_k H$ and whose multiplication is defined by

$$(a \# h)(b \# g) = \sum_{(h)} a(h_1 \cdot b) \# h_2 g,$$

where $\Delta(h) = \sum_{(h)} h_1 \otimes h_2$ is the comultiplication of h . Then $A \subseteq A \# H =: B$ is an extension with additional module structure whose module action is given by $a \# h \cdot b = a(h \cdot b)$. The subring of invariants is $A^B = A^H = \{a \in A \mid h \cdot a = \epsilon(h)a \ \forall h \in H\}$. For more details on Hopf algebra action we refer to [16].

4.1. Regularity of the subring of invariants. From 2.5 we get a characterisation of the regularity of the subring of invariants of A .

Proposition. *Let A be a k -algebra with Hopf action H . Then A^H is regular if and only if A is a semi-projective left $A \# H$ -module such that any cyclic left ideal generated by an H -invariant element is generated by an H -invariant idempotent.*

4.2. In order to ensure that A is a finitely presented object in $\sigma[A\#H A]$ we will assume some finiteness conditions on H or on its action. We say that a Hopf algebra H *acts finitely* on a k -algebra A if the image of the defining action $H \rightarrow \text{End}_k(A)$ is a finitely generated k -module or equivalently if $A\#H/\text{Ann}_{A\#H}(A)$ is finitely generated as left A -module. Recall that a k -algebra A is called *affine* if it is finitely generated as k -algebra.

Denote by $\epsilon : H \rightarrow k$ the counit of H . We need the following Lemma:

Lemma. *Let H be a Hopf algebra over k that is affine as k -algebra, then $\text{Ker}(\epsilon)$ is a finitely generated left ideal.*

Proof. Suppose that H is affine and let $\mathcal{B} \subseteq H$ be a finite set of elements which generate H as a k -algebra. We will show that $\text{Ker}(\epsilon) = \sum_{b \in \mathcal{B}} H(b - \epsilon(b))$. Obviously the right hand side is included in the left hand side. Note that for any word (=product) $\omega = b_1 \cdots b_m$ with $b_i \in \mathcal{B}$ we might set $a_0 = \epsilon(\omega)$, $a_m = \omega$ and $a_i = b_1 \cdots b_i \epsilon(b_{i+1} \cdots b_m)$ for $0 < i < m$ and conclude that $\omega - \epsilon(\omega) \in \sum_{i=1}^m H(b_i - \epsilon(b_i))$, since as a telescopic sum we have

$$\omega - \epsilon(\omega) = \sum_{i=1}^m a_i - a_{i-1} = \sum_{i=1}^m b_1 \cdots b_{i-1} \epsilon(b_{i+1} \cdots b_m) (b_i - \epsilon(b_i)).$$

Take any element $h \in \text{Ker}(\epsilon)$. Then there exist $\lambda_i \in k$ and words ω_i in $\mathcal{B}$ such that

$$h = h - \epsilon(h) = \sum_i \lambda_i [\omega_i - \epsilon(\omega_i)] \in \sum_{b \in \mathcal{B}} H(b - \epsilon(b)).$$

Thus $\text{Ker}(\epsilon)$ is finitely generated. $\square$

In the telescopic sum argument in the proof of the last Lemma we made use of the fact that the counit ϵ of a Hopf algebra is an algebra homomorphism. We do not know whether this Lemma holds true for affine weak Hopf algebras.

4.3. From Proposition 2.1 we deduce the next result:

Theorem. *If H is an affine k -algebra or acts finitely on A , then A is a finitely presented in $\sigma[A\#H A]$ and the following statements are equivalent:*

- (1) *A is H -regular, i.e. any finitely generated H -stable left ideal is generated by an H -invariant element;*
- (2) *A is a projective generator in $\sigma[A\#H A]$ and any cyclic left ideal generated by an H -invariant element is generated by an H -invariant idempotent.*
- (3) *A^H is von Neumann regular and $(-)^H : \sigma[A\#H A] \rightarrow A^H\text{-Mod}$ is an equivalence.*
- (4) *A^H is von Neumann regular and A is a projective generator in $\sigma[A\#H A]$.*
- (5) *A is a regular module in $\sigma[A\#H A]$.*

Proof. Once we showed that A is finitely presented in $\sigma[A\#H A]$, the result follows from 2.1. If H acts finitely on A , then we might substitute $A\#H$ by $B = A\#H/\text{Ann}_{A\#H}(A)$ which is finitely generated as left B -module. Also α lifts to a map $\bar{\alpha} : B \rightarrow A$ and splits as left A -module map. Thus $\text{Ker}(\bar{\alpha})$ is finitely generated as left A -module and thus as left ideal of B , i.e. A is finitely presented in $B\text{-Mod}$ and hence in $\sigma[B A] = \sigma[A\#H A]$.

On the other hand suppose that H is affine, then $\text{Ker}(\epsilon)$ is a finitely generated left ideal by Lemma 4.2. For the module algebra A and $\text{Ker}(\alpha : A\#H \rightarrow A)$, we have that if $x = \sum_{i=1}^n a_i \# h_i \in$

$\text{Ker}(\alpha)$, then

$$x = \sum_{i=1}^n a_i \# h_i - \left(\sum_{i=1}^n a_i \epsilon(h_i) \right) \# 1 = \sum_{i=1}^n a_i \# (h_i - \epsilon(h_i)) \in A \# \text{Ker}(\epsilon).$$

Thus $\text{Ker}(\alpha) = A \# \text{Ker}(\epsilon) = \sum_{b \in \mathcal{B}} A \# H(1 \# (b - \epsilon(b)))$ is a finitely generated left ideal of $A \# H$ and therefore A is finitely presented. $\square$

4.4. Note that the notion of regularity used here is different from the concept of an H -regular module algebra as defined by [26]. There the author define an element a of an H -module algebra A to be H -regular if $a \in (H \cdot a)A(H \cdot a)$ and calls A H -regular if every element is H -regular.

4.5. **The envelopping Hopf algebroid.** In general a Hopf action does not extend to the envelopping algebra A^e unless H is cocommutative. In order to study the two-sided H -stable ideals of a Hopf module algebra A with Hopf action H , one defines a new product on the tensor product $A^e \otimes H$ as follows:

$$[(a \otimes b) \bowtie h][(a' \otimes b') \bowtie h'] = \sum_{(h)} a(h_1 \cdot a') \otimes (h_3 \cdot b')b \bowtie h_2 h'$$

for all $a \otimes b, a' \otimes b' \in A^e$ and $h, h' \in H$. This construction had been used by the author in [14] (see also [13]) in order to define the central closure of a module algebra A as the self-injective hull of A as $A^e \bowtie H$ -module and had also been used by Connes and Moscovici in [7]). A similar construction had been used by L.Kadison in [11] which in [18] was shown to be isomorphic to the construction by Connes-Moscovici. Following Kadison, we denote this algebra on $A^e \otimes H$ by $A^e \bowtie H$ and call it the *envelopping Hopf algebroid* of A and H . For any left $A^e \bowtie H$ -module M denote by

$$Z(M)^H := \{m \in M \mid am = ma \wedge hm = \epsilon(h)m \forall a \in A, h \in H\}.$$

Then since $A \subseteq A^e \bowtie H$ is again an extension with additional module structure we have that $Z(-)^H$ is a functor from $A^e \bowtie H \rightarrow Z(A)^H\text{-Mod}$ and that

$$\text{Hom}_{A^e \bowtie H}(A, M) \rightarrow Z(M)^H \quad f \mapsto (1)f$$

is a functorial isomorphism. Note that $Z(A)^H := Z(A) \cap A^H \simeq \text{End}_{A^e \bowtie H}(A)$.

From 2.5 we get a characterisation of the regularity of the subring of central invariants of A .

Corollary. $Z(A)^H$ is regular if and only if A is a semi-projective left $A^e \bowtie H$ -module such that any cyclic ideal generated by a central H -invariant element is generated by a central H -invariant idempotent.

4.6. As before we need to ensure that A is a finitely presented object in $\sigma_{[A^e \bowtie H]A}$ in order to apply 2.1.

Lemma. If A and H are affine k -algebras, then A is a finitely presented module in $\sigma_{[A^e \bowtie H]A}$.

Proof. Consider $\alpha : A^e \bowtie H \rightarrow A$ by $a \otimes b \bowtie h \mapsto a\epsilon(h)b$. For any $x = \sum_{i=1}^n a_i \otimes b_i \bowtie h_i \in \text{Ker}(\alpha)$ we have $\sum_{i=1}^n a_i \epsilon(h_i) b_i = 0$. Hence

$$\begin{aligned} x &= \sum_{i=1}^n a_i \otimes b_i \bowtie h_i - \left(\sum_{i=1}^n a_i b_i \epsilon(h_i) \right) \otimes 1 \bowtie 1 + \left[\sum_{i=1}^n a_i \otimes b_i \bowtie \epsilon(h_i) - \sum_{i=1}^n a_i \otimes b_i \bowtie \epsilon(h_i) \right] \\ &= \sum_{i=1}^n a_i \otimes b_i \bowtie [h_i - \epsilon(h_i)] + \sum_{i=1}^n a_i \epsilon(h_i) [1 \otimes b_i - b_i \otimes 1] \bowtie 1 \\ &\in A^e \bowtie \text{Ker}(\epsilon) + A \text{Ker}(\mu) \otimes 1 \end{aligned}$$

where $\text{Ker}(\mu)$ is the augmentation ideal of the envelopping algebra, i.e. the kernel of the multiplication map $\mu : A^e \rightarrow A$. Hence we see that $\text{Ker}(\alpha)$ is generated as left ideal of $A^e \bowtie H$ by elements of $1 \otimes \text{Ker}(\epsilon)$ and $\text{Ker}(\mu) \bowtie 1$. It is well-known that $\text{Ker}(\mu)$ is finitely generated left ideal of A^e if A is affine and by 4.2 it follows that $\text{Ker}(\epsilon)$ is finitely generated if H is affine. $\square$

4.7. H -biregular module algebras. The last statement, 2.1 and 2.4 yield the main result in this section which generalises [4, 1.2] from group actions to Hopf actions.

Corollary. *Let A and H be affine k -algebras, then the following statements are equivalent:*

- (a) *A is H -biregular, i.e. every finitely generated H -stable two-sided ideal of A is generated by a central H -invariant idempotent.*
- (b) *A is a projective generator in $\sigma[A^e \bowtie H A]$ and any ideal generated by a central H -invariant element is generated by an idempotent central H -invariant element.*
- (c) *A is a regular module in $\sigma[A^e \bowtie H A]$.*
- (d) *$Z(A)^H$ is von Neumann regular and one of the following statements hold:*
 - (i) *the functor $Z(-)^H : \sigma[A^e \bowtie H A] \rightarrow Z(A)^H\text{-Mod}$ is an equivalence.*
 - (ii) *A is a projective generator in $\sigma[A^e \bowtie H A]$.*
 - (iii) *every maximal H -stable ideal M of A can be written as $M = [Z(A)^H \cap M]A$.*
 - (iv) *A_m is H -simple for any maximal ideal m of $Z(A)^H$.*

4.8. Relative semisimple extension. Let G be a finite group acting on an algebra A . The condition that $|G|$ is invertible is frequently used in the study of group actions because it implies that $A \subseteq A * G$ is a separable extension. The weaker condition on A of having an *element of trace 1*, i.e. an element $z \in A$ such that $t \cdot a = \sum_{g \in G} (g \cdot a) = 1$, where $t = \sum_{g \in G} g$, implies at least the projectivity of A as $A * G$ -module. Here we will analyse those concepts and carry them over to Hopf algebra actions.

The antipode of a Hopf algebra H is denoted by S . An element $t \in H$ is called a right (resp. left) integral in H if $th = t\epsilon(h)$ (resp. $ht = \epsilon(h)t$) for all $h \in H$. It is well-known that $\sum_{(t)} S(t_1) \otimes t_2 h = \sum_{(t)} h S(t_1) \otimes t_2$ for all $h \in H$.

Proposition. *Let H be a Hopf algebra and A a left H -module algebra. Suppose H has a non-zero right integral t and A admits a central element z such that $S(t) \cdot z = 1$, then*

$$c := \sum_{(t)} (1 \# S(t_1)) \otimes (z \# t_2)$$

is a Casimir element of $A \# H$ that acts unitarily on A . Hence A is a semisimple $(A \# H, A)$ -module.

- (1) *$A \subseteq A \# H$ is a semisimple extension if $A^H \subseteq A$ is a H^* -Galois extension, i.e. A is a generator in $A \# H\text{-Mod}$.*
- (2) *$A \subseteq A \# H$ is separable if $z \in A^H$ or H is cocommutative*

Proof. The element $\sum_{(t)} 1 \# S(t_1) \otimes z \# t_2$ is a Casimir element in $(A \# H) \otimes_A (A \# H)$ because for all $a \# h \in A \# H$:

$$\begin{aligned}
c(a\#h) &= \sum_{(t)} (1\#S(t_1)) \otimes (z\#t_2)(a\#h) \\
&= \sum_{(t)} (1\#S(t_1)) \otimes z(t_2 \cdot a)\#t_3h \\
&= \sum_{(t)} (S(t_2) \cdot (t_3 \cdot a))\#S(t_1) \otimes z\#t_4h \\
&= (a\#1) \left(\sum_{(t)} 1\#S(t_1) \otimes z\#t_2h \right) \\
&= (a\#1) \left(\sum_{(t)} 1\#hS(t_1) \otimes z\#t_2 \right) \\
&= (a\#h) \left(\sum_{(t)} 1\#hS(t_1) \otimes z\#t_2 \right) = (a\#h)c
\end{aligned}$$

Moreover

$$\sum_{(t)} (1\#S(t_1))(z\#t_2) \cdot 1 = \sum_{(t)} S(t_1) \cdot (z(t_2 \cdot 1)) = S(t) \cdot z = 1.$$

Thus c acts unitarily on A . By 3, A is a semisimple $(A\#H, A)$ -module.

If A/A^H is a H^* -Galois extension, then $\sigma[A\#H A] = A\#H$ and the claim follows from 3.2.

Note that $\mu(c) = \sum_{(t)} S(t_2)z\#S(t_1)t_3$. If $z \in Z(A)^H$, then $\mu(c) = \sum_{(t)} z\#S(t_1)\epsilon(t_2)t_3 = \epsilon(S(t))z\#1 = S(t)z\#1 = 1\#1$. If H is cocommutative, then $\mu(c) = \sum_{(t)} S(t_1)z\#S(t_2)t_3 = S(t)x\#1 = 1\#1$. Hence in both cases $A\#H$ is separable over A . $\square$

4.9. If the antipode is bijective and A has a central element of trace 1, i.e. $z \in Z(A)$ with $t \cdot z = 1$ for a left integral t of H , then $t' = S^{-1}(t)$ is a right integral, and $S(t') \cdot z = 1$ holds, i.e. the condition of 4.8 is fulfilled.

4.10. The observation that $A \subseteq A\#H$ is separable if $z \in Z(A)^H$ or H cocommutative, can also be found in [5, Theorem 1.11] or [6], but under the hypothesis of H being a Frobenius k -algebra and thus finitely generated and projective as k -module. Note that the existence of a left (or right) integral forces a Hopf algebra in many cases to be finitely generated although there are examples of non-finitely generated ones (see [12]).

4.11. **Regularity of smash products.** In [1] the authors studied the regularity of skew group rings. They showed in particular that a skew group ring $A * G$ is regular if A is regular, G is locally finite and for every finite subgroup H of G there exists a central element of H -trace 1. In this section we will show how much of their arguments go over to smash products.

4.12. First note the following Corollary that we get from 4.8:

Corollary. *Let H be a Hopf algebra acting on a regular module algebra A . Assume that there exists a right integral t of H and a central element z such that $S(t) \cdot z = 1$.*

- (1) *If H acts finitely on A , then A is regular in $\sigma[A\#H A]$, A^H is regular and $A^H\text{-Mod}$ is Morita-equivalent to $\sigma[A\#H A]$.*
- (2) *If z is H -invariant or H is cocommutative or A/A^H is H^* -Galois and ${}_k H$ is finitely generated, then $A\#H$ is regular.*

Proof. (1) If we substitute $A\#H$ by $B = A\#H/\text{Ann}(A)$, then ${}_AB$ is finitely generated and $c = \sum_{(t)} 1\#S(t_1) \otimes z\#t_2 \in A\#H \otimes_A A\#H$ can be lifted to $c' \in B \otimes_A B$ which still acts unitarily on A . By Corollary 3.2, A has the properties stated above.

(2) if $z \in A^H$ or H is cocommutative, then by 4.8, $A \subseteq A\#H$ is separable and hence $A\#H$ is regular as A was regular. In case A/A^H is H^* -Galois we have that $\sigma_{[A\#H A]} = A\#H\text{-Mod}$ and by 4.8, $A \subseteq A\#H$ is a semisimple extension. Since ${}_kH$ is finitely generated, ${}_AA\#H$ is finitely generated. Hence any cyclic left ideal I of $A\#H$ is finitely generated as left A -submodule of $A\#H$. Since A is regular I is a direct summand of $A\#H$ and as $A \subseteq A\#H$ is semisimple, it is also a direct summand of $A\#H$ as left ideal. $\square$

4.13. Locally finite Hopf algebras. Call an extension $A \subseteq B$ *locally separable* if every element of B is contained in an intermediate algebra $A \subseteq C \subseteq B$ such that C is a separable extension of A (see also A. Magid's definition [15]). Of course, if $A \subseteq B$ is locally separable and A is regular, then B is regular, because any element $x \in B$ is contained in a separable extension C of A . And if A is regular, then also C . Thus $x = xyx$ for some $y \in C \subseteq B$. Hence B is regular. The characterisation of regular group rings $k[G]$ can actually be stated as $k \subseteq k[G]$ being locally separable and k being regular. Alfaro, Ara and del Rio proved in [1, Theorem 1.3] that if G is a locally finite group acting on a regular ring A such that for every finite subgroup H there exists a central element of trace 1 with respect to H , then the skew group ring $A * G$ is regular. We will slightly generalize there result to Hopf algebra actions by showing that the hypotheses of their result imply that $A \subseteq A * G$ is locally separable.

A group is called locally finite if any finitely generated subgroup is finite.

Definition. Let H be a Hopf algebra over k . A Hopf algebra is called *locally finite* if any finite set $X \subseteq H$ is contained in a Hopf subalgebra of H which contains a non-zero right integral.

Note that any Hopf algebra H which is free as a module over k is finitely generated as k -module if and only if it contains a non-zero right integral (see [12]). A group ring $k[G]$ is of course locally finite if G is locally finite.

4.14. We are now in position to generalize [1, Theorem 1.3]:

Corollary. Let H be a locally finite Hopf algebra over k acting on a k -algebra A such that for any Hopf subalgebra K of H that contains a non-zero right integral t , there exists a central element $z_t \in A$ with $S(t) \cdot z_t = 1$. If H is cocommutative or $z_t \in Z(A)^H$ for all right integrals t , then $A \subseteq A\#H$ is locally separable. Hence if A is regular, so is $A\#H$.

Proof. Let $x := \sum_i^k a_i \# h_i \in A\#H$. By hypothesis $K := \langle \{h_1, \dots, h_k\} \rangle$ contains a non-zero right integral t . By 4.8(2), $A \subseteq A\#K$ is separable, and hence regular if A was regular. $\square$

4.15. As a consequence we have that if H is a cocommutative Hopf algebra acting on a commutative regular k -algebra having a central element of trace 1, then $A\#H$ is regular (which partly generalizes [1, Corollary 2.5]). It had been shown in [1, 2.4], that if a skew-group ring $A * G$ is regular, then is also A . This is not anymore true for smash products as it is easily seen by the fact, that for any finite dimensional Hopf algebra H over a field k , the smash product $H\#H^* \simeq M_n(k)$ is isomorphic to a semisimple artinian ring, whether H is semisimple or not. However we have that if H is an n -dimensional cosemisimple Hopf algebra over a field k acting on an algebra A such that $A\#H$ is regular, then A is regular. Simply because by the Blattner-Montgomery duality one

has $(A\#H)\#H^* \simeq M_n(A)$ and since H^* is separable over k , we have $A\#H\#H^*$ being separable over $A\#H$ inducing regularity on $M_n(A)$ and hence on A .

5. REGULARITY AND INJECTIVITY OF THE SUBRING OF INVARIANTS

Note that from Zelmanowitz result 2.6 we get

Corollary. *Let A be an H -module algebra that is projective in $\sigma[A]$. If A^H is large in A then A^H is left self-injective and von Neumann regular if and only if A is a self-injective left $A\#H$ -module which is non-singular in $\sigma[A\#H]$. In this case A is also H -semiprime.*

Proof. The equivalence of the statements follows verbatim from 2.6. If I is an H -stable ideal of A with $I^2 = 0$, then $(I^H)^2 = 0$. But since A^H is regular, $I^H = 0$ and since A^H is large $I = 0$. $\square$

5.1. To compare the injectivity of A and its subring of invariants, we need the following Lemma which is probably known:

Lemma. *Let $S \subseteq T$ be rings such that T_S is flat. If T is left self-injective, then so is S .*

Proof. Let I be a left ideal of S , denote the inclusion map by $f : I \rightarrow S$ and let $g : I \rightarrow S$ be an S -linear map. Let $\gamma : T \rightarrow T \otimes_S S$ be the canonical isomorphism. Then γ is left T -linear and $\gamma|_{TI} : TI \rightarrow T \otimes_S I$ is also an isomorphism of left T -modules. Let $\tilde{f} := \gamma(1 \otimes f)\gamma^{-1} : TI \rightarrow T$. As T_S is flat, $\tilde{f}$ is injective. Also set $\tilde{g} := \gamma(1 \otimes g)\gamma^{-1} : TI \rightarrow T$. Then we can consider the following diagram with exact rows, where $\iota : S \rightarrow T$ denotes the inclusion map (which is of course just S -linear):

$$\begin{array}{ccccc} 0 & \longrightarrow & I & \xrightarrow{f} & S \\ & & \downarrow \iota & & \downarrow \iota \\ 0 & \longrightarrow & TI & \xrightarrow{\tilde{f}} & T \\ & & \downarrow \tilde{g} & & \\ & & T & & \end{array}$$

As T is left self-injective, there exists a T -linear map $\tilde{h} : T \rightarrow T$ such that $\tilde{f}\tilde{h} = \tilde{g}$. Hence the outer trapezoid also commutes, i.e. $f\iota\tilde{h} = \iota\tilde{g}$. Since for all $x \in I$: $(x)\iota\tilde{g} = (x)g$ we may identify $\iota\tilde{g}$ with g and take $h := \iota\tilde{h}$ to be the desired S -linear map. $\square$

5.2. We will finish with the following result on the transfer of regularity and injectivity to the subring of invariants of a module algebra which should be compared to [9, Theorem A].

Corollary. *Let H be Hopf algebra acting on A . Suppose H has a right integral t and A has a central element z such that $S(t) \cdot z = 1$. If A is regular and left self-injective ring, then A^H is regular and left self-injective.*

Proof. Since A is $(A\#H.A)$ -semisimple by 4.8, A is semi-projective as $A\#H$ -module. Take any $x \in A^H$, then Ax is a direct summand in A and by relative semisimplicity also a direct summand as $A\#H$ -submodule. Thus by 5.1 A^H regular. Now it follows from (1) that A^H is also left self-injective. $\square$

REFERENCES

- [1] R. Alfaro, P.Ara and Angel del Rio, *Regular skew group rings.*, J.Austral. Math. Soc. (Series A) 58(1995), 167-182
- [2] M.Cabrera, and A.A.Mohammed, *Extended centroid and central closure of multiplicatively semiprime algebras.*, Comm. Algebra **29** (2001), no. 3, 1215-1233
- [3] S.Caenepeel, T. Guédénon, *Fully bounded Noetherian rings and Frobenius extensions.*, Journal of Algebra and its Application 6, No. 2, 189-206 (2007).
- [4] Castella, D., *Birégularité des anneaux de groupes gauches et des extensions libérales.*, Commun. Algebra 19, No.10, 2787-2811 (1991).
- [5] M.Cohen, D.Fishman, *Semisimple extensions and Elements of Trace 1*, J. Alg. 149, 419-437 (1992)
- [6] M.Cohen, D.Fishman, *Erratum "Semisimple extensions and Elements of Trace 1"*, J. Alg. 165, 242 (1994)
- [7] A. Connes, H. Moscovici, *Rankin-Cohen brackets and the Hopf algebra of transverse geometry*, Moscow Math. J. 4 (2004), 111-130.
- [8] D.Fieldhouse, *Regular rings and modules.*, J. Aust. Math. Soc. 13, 477-491 (1972).
- [9] J.M.Goursad, J.Osterburg, J.L.Pascaud, J.Valette, *Points fixes des anneaux réguliers auto-injectifs à gauche*, Comm.Alg. (1981).
- [10] K.Hirata, K.Sugano, *On semisimple extensions and separable extensions over non commutative rings.*, J. Math. Soc. Japan, 18(4):360-373, 1966.
- [11] L.Kadison, *Pseudo-Galois Extensions and Hopf Algebroids in Modules and Comodules*, eds. T.Brzezinski, J.L. Gómez Pardo, I.Shestakov, P.Smith, Trends in Mathematics (Birkhäuser) 247-264 (2008)
- [12] C.Lomp, *Integrals in Hopf algebras over rings*, Communications in Algebra, 32(12), 4687-4711 (2004)
- [13] C.Lomp, *A central closure construction for certain extensions. Applications to Hopf actions*, Journal of Pure and applied Algebra, 198 (1-3), 297-316 (2005)
- [14] C.Lomp, *Primeigenschaften von Algebren in Modulkategorien über Hopfalgebren*, PhD Thesis, Heinrich-Heine Universität Düsseldorf (2002)
- [15] A.R. Magid, *Locally Galois algebras.*, Pacific Journal of Mathematics. Volume 33, Number 3 (1970), 707-724.
- [16] S. Montgomery. *Hopf Algebras and Their Actions on Rings.*, CBMS Regional Conference Series in Mathematics 82, AMS (1992)
- [17] J. von Neumann, *Regular rings*, Proc. Nat. Acad.Sci. USA 22(1936),707-713
- [18] F.Panaite, F.Van Oystaeyen, *Some Bialgebroids Constructed by Kadison and Connes-Moscovici are Isomorphic*, Applied Categorical Structures 14(5-6), 627-632 (2006)
- [19] P.Schauenburg, *Weak Hopf algebras and quantum groupoids*, Banach Center Publ. 61 (2003), 171-188
- [20] A.Tuganbaev, *Semidistributive Modules and Rings*, Kluwer Academics Publishers, Dordrecht (1998)
- [21] R.Wisbauer, *Nonassociative left regular and biregular rings*, J. Pure Appl Algebra 10 (1977), 215-226
- [22] R.Wisbauer, *Foundations of Module and Ring Theory.*, Gordon and Breach (1991)
- [23] R.Wisbauer, *Modules and algebras: bimodule structure and group actions on algebras.*, Addison Wesley Longman Ltd. 1996
- [24] J.Zelmanowitz, *Regular modules*, Trans. Amer. Math. Soc. 163 (1972), 341-355.
- [25] J.Zelmanowitz, *Density for polyform modules.*, Huynh, D. V. (ed.) et al., *Algebra and its applications.*, Contemp. Math. 259, 563-569 (2000).
- [26] S.Zhang, *The Von Neumann Regular Radical and Jacobson Radical of Crossed Products*, Acta Mathematica Hungarica, Volume 86, Number 4, March 2000, 319-333(15)

UNIVERSITY OF PORTO, DEPARTMENT OF PURE MATHEMATICS, RUA CAMPO ALEGRE 687, 4169-007 PORTO (PORTUGAL), clomp@fc.up.pt